

Levomepromazine

Antipsychotic



Your medication

Levomepromazine



What is this leaflet for?

This leaflet is to help you understand more about your medicine.

Your medicine could look different to the pictures on this leaflet.



Levomepromazine is also called Nozinan.

Levomepromazine is available as tablets and injection.




What is Levomepromazine used for?

Levomepromazine is commonly used to treat psychosis.

It is also used in people having behavioural problems.



Levomepromazine



What will Levomepromazine do?

Levomepromazine should help you feel calm and relaxed.

It could take some time for Levomepromazine to have its full effect.



What are the side effects of Levomepromazine?

Side effects are the unwanted effects of any medicine on the body.

Not everyone taking a medicine will get side effects.

Many side effects will go away with time. But some side effects may last for longer.

Some side effects can be serious.

You should tell your doctor if you have any side effect.

Levomepromazine



What are the side effects of Levomepromazine?

- feeling dizzy
- feeling restless or cannot sit still
- stiffness in arms or legs
- feeling sleepy or sluggish
- increase in weight
- changes in blood sugar level
- changes in cholesterol levels
- changes in heart function
- rash or feeling itchy
- in women, it can effect periods.



You will need to have the following tests done as part of the monitoring for side effects:

Tracing of your heart - ECG


Blood pressure

Weight

Waist circumference

Blood tests

Levomepromazine



How and when to take Levomepromazine?

The label on your Levomepromazine packet should tell you when and how much you should take.



What else can help?

Eating healthy

Doing some exercises.



Participating in activities that you like.



If you forget to take the tablet


Tell your carer or doctor straight away.



When to stop taking Levomepromazine?

You should carry on taking your medicine unless your doctor asks you to stop.

Levomepromazine



Some medicines do not work well together.

You should tell your doctor about any other medicines you are taking.



What should not be done while taking Levomepromazine?

Drinking alcohol when on Levomepromazine could make you feel drowsy.



Doctor's comments

Help and support



Your doctor:
Telephone Number:



Your Psychiatrist: Dr S Bhaumik
Telephone Number:



Your Community Nurse:
Telephone Number:

Other professionals involved:

Patient  UK
The same health information as provided
by GPs to patients during consultations

Other resources:

www.patient.co.uk

NHS CALL 24 HOURS A DAY
0845
Direct 4647



Leaflet produced and issued by
Leicestershire Partnership NHS Trust
www.leicspt.nhs.uk

Pictures used from Change Picture Bank and Somerset Symbols Project

Thanks to G. Unwin and S. Deb from Birmingham University for use of the content of their medication leaflets.

Date implemented: May 2011

Review Date: May 2013

Leaflet no: 189 – Edition 1

Please note this is not a manufacturer's information leaflet